



SEED SOWING INSTRUCTIONS

WAYFARING

Treating and growing on seed

Need: Seed, cleaning equipment [see 2.a], polythene freezer bags, vermiculite [available from Garden Centres], seed tray/rigipot, peat free compost, sharp sand, liquid fertiliser. Optional: fine grit, watering tray.

1. STORAGE

Clean flesh off seed.

Technique: sieve / rolling pin / food blender are useful tools.

Wash seed thoroughly to remove all flesh [which can contain inhibitors to germination], and store in plastic bag at 10- 15° C until late January- tie top loosely.

Allow seed to breathe by opening/ shaking every 2/ 3 weeks.

2. STRATIFICATION

In early October start process. Steps:

a] *Preparation*

Soak seed for 24- 48 hours to remoisturise.

Mix vermiculite and compost with seed at ratio of 3 parts seed to one part compost, and place in polythene freezer bag or use suitable plastic container if rodents are a problem.

Add water until damp - not much required - use a beaker or similar container.. Labelseed inside bag and tie off top of bag.

Mark up bag with seed collection details : species, collection site, date.

b] *Pretreatment*

Store in cold conditions [2°-5° C] for 8 weeks **or** as long as seed takes to 'chit' [produce a root] - use fridge.

Once a week open bag : mix seeds allowing carbon dioxide to escape and oxygen in. Re-moisten if necessary and remove any mouldy seeds.

At first signs of chitting, check bag regularly for root emergence.

3. **SOWING**

Fill seed tray/rigipot cells with compost to 1-2 cms. from top of tray. Bang on table to firm.

Sow seed in growing tray when root is about 1/2" long, using peat free sowing / potting compost .

Fill tray asap with chitted seed [keeps growth even].

Do not sow seed with too long a shoot in a shallow seed tray. Sow seeds with long root in rigipot which has deep cells.

Cover compost in cells with fine grit / coarse sand if available.

4. **GROWING**

a] Positioning of tray

Put trays of early seedlings on windowsill/in **warm** greenhouse to avoid winter cold/frosts; if available move into cold frame/greenhouse in April/May, or else leave on windowsill.

Move trees outside in September to overwinter, avoiding excessive sunlight.

Ensure slugs / snails / rodents, etc. cannot get at the seedlings.

Turn the tray regularly to avoid bending of tree stems [seedlings grow towards the light].

b] Watering:

Water regularly- every 3-4 days in mid-summer. Water less regularly in winter, but enough to keep roots damp.

It is difficult to overwater in summer - if in doubt, be generous.

c] Feeding

Mix in liquid fertiliser when watering, if necessary. Ask for assistance if unclear.

6. ***ADVICE SERVICE AND HELPLINE***

Please telephone COLIN CARPENTER if you have any queries, on :
01525 716371 OR MOBILE 07751 356094